arXiv:1205.5715v1 [gr-qc] 25 May 2012

KEK-TH-1550

Time evolution of a thin black ring via Hawking radiation

Mitsuhiro Matsumoto(1), Hirotaka Yoshino(2), and Hideo Kodama(1,2)
(1) *Department of Particle and Nuclear Physics,*
The Graduate University for Advanced Studies (SOKENDAI),
Tsukuba 305-0801, Japan and
(2) *Theory Center, Institute of Particles and Nuclear Studies,*
KEK, Tsukuba, Ibaraki, 305-0801, Japan

(Dated: May 21, 2012)

Abstract

Black objects lose their mass and angular momenta by Hawking radiation. It was shown that if only scalar fields are emitted, a five-dimensional Myers-Perry black hole evolves toward a state with nonvanishing nondimensional rotation parameters, similarly to the case of a four-dimensional Kerr black hole. In this paper, we extend this analysis to the case of a black ring. By approximating a thin Pomeransky-Sen'kov black ring by a boosted Kerr string, we derive a set of equations that determines time evolution of quasistationary evaporation of the black ring. Our result shows that a thin black ring evaporates without changing the nondimensional rotational parameter along S^1. In the case of an Emparan-Reall black ring, we solve those equations semianalytically and determine its time evolution.

I. INTRODUCTION

In four spacetime dimensions, a stationary, asymptotically flat, vacuum black hole is completely characterized by its mass and spin [1]. In particular, the topology of its event horizon must be a sphere [2]. By contrast, in five dimensions, in addition to the Myers-Perry black hole [3] which is natural generalization of the four-dimensional Kerr black hole, various exact solutions of black objects with nonspherical horizon topologies have been found (see [4] for a review). In this paper, we focus attention to the black ring solution with the $S^1 \times S^2$ horizon topology. The black ring solution rotating in the direction of S^1 was found by Emparan and Reall [5]. Since a five-dimensional spacetime can have two angular momenta, Pomeransky and Sen'kov [6] extended it to black rings with two independent rotation parameters (i.e., doubly spinning in directions of S^1 and S^2).

A black hole is known to evaporate due to the effect of quantum field theory in curved spacetime as shown by Hawking [7]. The rate of mass and angular momentum loss of a Kerr black hole by the Hawking radiation was first studied by Page [8, 9] taking account of fields with spins 1/2, 1, and 2, and it was shown that a Kerr black hole spins down to a nonrotating black hole regardless of its initial state. However, Chambers *et al.* [10] (see also [11]) showed that if only massless scalar fields are taken into account (i.e., in the absence of fields with nonzero spin), a four-dimensional Kerr black hole evolves to a state with the nonvanishing nondimensional rotation parameter, $a/M \simeq 0.555$. This analysis was extended to five-dimensional Myers-Perry black holes by Nomura *et al* [12]. They showed that any such black hole with nonzero rotation parameters a and b evolves toward an asymptotic state with $a/M^{1/2} = b/M^{1/2} \simeq 0.1975(8/3\pi)^{1/2}$. Here, this value is independent of the initial values of a and b.

It is interesting to extend these studies to the case of a black ring. Although the Hawking radiation of black rings has been studied in various context [13–16], so far, the time evolution of a black ring has not been studied. The difficulty in this study is that the method of mode decomposition of the Klein-Gordon field in this spacetime is not known since separation of variables has not been done, and therefore, two dimensional numerical calculations of eigenfunctions are required. In order to avoid this difficulty, we take the thin limit and approximate a black ring as a boosted black string. In this approximation, the variables can be separated and we have well defined modes.

Using this approximation, we give a formulation to study the evolution of a thin Pomeransky-Sen'kov black ring by the Hawking radiation, and discuss general features that do not depend on details of greybody factors. Then, we derive a semianalytic solution in the case of an Emparan-Reall black ring and determine its time evolution. Here, the solution is semianalytic in the sense that one parameter that is related the greybody factors has to be evaluated numerically, although the evolution is expressed by analytic formulas.

This paper is organized as follows. In Sec. II, the black ring solution is reviewed and its boosted Kerr string limit is shown. In Sec. III, we derive the equations that determine the emission rates of mass and angular momenta of a black ring via Hawking radiation. In Sec. IV, the evolution of a black ring is discussed. Sec. V is devoted to a summary. In Appendix A, we check the validity of our numerical result by studying the DeWitt approximation. To simplify the notation, the natural units $\hbar = c = G = k_B = 1$ are used.

II. BLACK RING

In this section, we review basic properties of a black ring and show its boosted Kerr string limit. This limit was discussed in the more general cases of an unbalanced Pomeransky-Sen'kov black ring in Ref. [17].

The metric of the Pomeransky-Sen'kov black ring is [6]

$$\begin{aligned} ds^2 = &- \frac{H(y,x)}{H(x,y)} (dt + \Omega)^2 - \frac{F(x,y)}{H(y,x)} d\psi^2 - 2\frac{J(x,y)}{H(y,x)} d\psi d\phi + \frac{F(y,x)}{H(y,x)} d\phi^2 \\ &+ \frac{2R^2 H(x,y)}{(x-y)^2 (1-\nu)^2} \left(\frac{dx^2}{G(x)} - \frac{dy^2}{G(y)} \right), \end{aligned} \tag{1}$$

where the 1-form Ω is

$$\begin{aligned} \Omega = &- \frac{2R\lambda\sqrt{(1+\nu)^2 - \lambda^2}}{H(y,x)} \Big[\left(1 - x^2\right) y \sqrt{\nu} d\phi \\ &+ \frac{1+y}{1-\lambda+\nu} \left\{1 + \lambda - \nu + x^2 y \nu (1 - \lambda - \nu) + 2\nu x (1-y)\right\} d\psi \Big], \end{aligned} \tag{2}$$

and the functions G, H, J and F are

$$G(x) = \left(1 - x^2\right)\left(1 + \lambda x + \nu x^2\right), \tag{3}$$

$$H(x,y) = 1 + \lambda^2 - \nu^2 + 2\lambda\nu\left(1 - x^2\right) y + 2x\lambda\left(1 - y^2\nu^2\right) + x^2 y^2 \nu\left(1 - \lambda^2 - \nu^2\right), \tag{4}$$

$$J(x,y) = \frac{2R^2\left(1-x^2\right)\left(1-y^2\right)\lambda\sqrt{\nu}}{\left(x-y\right)\left(1-\nu\right)^2}\left[1+\lambda^2-\nu^2+2\left(x+y\right)\lambda\nu - xy\nu\left(1-\lambda^2-\nu^2\right)\right], \tag{5}$$

$$\begin{aligned} F(x,y) = & \frac{2R^2}{\left(x-y\right)\left(1-\nu\right)^2} \\ & \times \Bigg[G(x)\left(1-y^2\right)\left[\left\{(1-\nu)^2-\lambda^2\right\}(1+\nu)+y\lambda\left(1-\lambda^2+2\nu-3\nu^2\right)\right] \\ & \quad + G(y)\big[2\lambda^2+x\lambda\left\{(1-\nu)^2+\lambda^2\right\}+x^2\left\{(1-\nu)^2-\lambda^2\right\}(1+\nu) \\ & \qquad + x^3\lambda\left(1-\lambda^2-3\nu^2+2\nu^3\right)-x^4(1-\nu)\nu\left(-1+\lambda^2+\nu^2\right)\big]\Bigg]. \end{aligned} \tag{6}$$

Here, we follow the notation of Ref. [6] except that we choose the signature $(-,+,+,+,+)$ for the metric, exchange ϕ and ψ, and use R instead of k. The coordinate ranges are $-\infty < t < +\infty$, $0 < \phi, \psi < 2\pi$, $-1 \leq x \leq 1$ and $-\infty < y < -1$. R is a parameter of dimension of length, which determines the characteristic scale. λ and ν are dimensionless parameters satisfying $0 \leq \nu < 1$ and $2\sqrt{\nu} \leq \lambda < 1+\nu$, which determine two nondimensional rotational parameters. The regular event horizon exists at $y = y_h$, where

$$y_h = \frac{-\lambda + \sqrt{\lambda^2 - 4\nu}}{2\nu}. \tag{7}$$

The solution is asymptotically flat and spacelike infinity is located at $x = y = -1$. The coordinates (x, ϕ) parametrize the two-sphere S^2 and ψ parametrizes the circle S^1. One recovers the Emparan-Reall black ring by setting $\nu = 0$, and the line $\nu = \lambda^2/4$ represents the sequence of an extremal black ring (see Fig. 1).

The mass and angular momenta are

$$M = \frac{3\pi R^2 \lambda}{1-\lambda+\nu}, \qquad J_\phi = \frac{4\pi R^3 \lambda\sqrt{\nu}\sqrt{(1+\nu)^2-\lambda^2}}{(1-\nu)^2(1-\lambda+\nu)}, \tag{8}$$

$$J_\psi = \frac{2\pi R^3 \lambda(1+\lambda-6\nu+\lambda\nu+\nu^2)\sqrt{(1+\nu)^2-\lambda^2}}{(1-\nu)^2(1-\lambda+\nu)^2}. \tag{9}$$

The angular velocities, the area, and the surface gravity of the horizon are written as [18]

$$\Omega_\phi = \frac{\lambda(1+\nu)-(1-\nu)\sqrt{\lambda^2-4\nu}}{4R\lambda\sqrt{\nu}}\sqrt{\frac{1+\nu-\lambda}{1+\nu+\lambda}}, \qquad \Omega_\psi = \frac{1}{2R}\sqrt{\frac{1+\nu-\lambda}{1+\nu+\lambda}}, \tag{10}$$

$$A_H = \frac{32\pi^2 R^3 \lambda(1+\nu+\lambda)}{(1-\nu)^2(y_h^{-1}-y_h)}, \qquad \kappa = \frac{(y_h^{-1}-y_h)(1-\nu)\sqrt{\lambda^2-4\nu}}{4R\lambda(1+\nu+\lambda)}. \tag{11}$$

We consider a thin ring limit $\lambda \to 0$ where the ratio of the S^2 radius to the S^1 radius becomes very small. Here, we have to take care of the fact that this limit depends on the



FIG. 1: The parameter space (λ, ν) of the Pomeransky-Sen'kov solution. λ and ν can take values in the region surrounded by solid lines. The line $\nu = \lambda - 1$ is the Myers-Perry black hole limit, the line $\nu = 0$ is the Emparan-Reall black ring, the line $\nu = \lambda^2/4$ is the extremal limit. The point $\lambda = \nu = 0$ corresponds to the boosted Kerr string limit. The dashed line $\nu = a_*^2\lambda^2/4$ is the path to the point $\lambda = \nu = 0$ with a fixed a_*.

path to the point $\lambda = \nu = 0$. For example, taking limit $\lambda \to 0$ on the line $\nu = 0$ gives the boosted Schwarzschild string, while taking limit $\lambda \to 0$ on the extremal line $\nu = \lambda^2/4$ should result in a state different from the boosted Schwarzschild string. Therefore, $\lambda = \nu = 0$ is a degenerate point, and in order to resolve this degeneracy, we introduce a new parameter a_* as

$$\nu = \frac{1}{4} a_*^2 \lambda^2, \tag{12}$$

and consider the limit $\lambda \to 0$ on the line of a fixed a_* (see Fig. 1). Also, in order to obtain a well-defined limit, we introduce

$$M_K = \frac{1}{\sqrt{2}} \lambda R, \tag{13}$$

and fix M_K in taking this limit.

We introduce the new coordinates r, z and θ by

$$y = -\frac{\sqrt{2}R}{r}, \qquad \psi = -\frac{z}{\sqrt{2}R}, \qquad x = \cos\theta, \tag{14}$$

and collect the leading-order term of each metric component with respect to λ. Then, the black ring solution is reduced to the so-called boosted Kerr string solution

$$\begin{aligned}ds^2 = &-\left(1-\frac{2M_K r\cosh^2\sigma}{\rho^2}\right)dt^2+\frac{2M_K r\sinh 2\sigma}{\rho^2}dtdz+\left(1+\frac{2M_K r\sinh^2\sigma}{\rho^2}\right)dz^2\\ &+\frac{\rho^2}{\Delta}dr^2+\rho^2 d\theta^2+\frac{\left(r^2+a^2\right)^2-\Delta a^2\sin^2\theta}{\rho^2}\sin^2\theta d\phi^2\\ &-\frac{4M_K r\cosh\sigma}{\rho^2}a\sin^2\theta dtd\phi-\frac{4M_K r\sinh\sigma}{\rho^2}a\sin^2\theta dzd\phi,\end{aligned}\tag{15}$$

where $\rho^2=r^2+a^2\cos^2\theta$ and $\Delta=r^2-2M_K r+a^2$, and a is defined by $a:=M_K a_*$. Since M_K is the mass density and a is the Kerr parameter, a_* represents the nondimensional rotation parameter along S^2 direction. $\sigma:=\mathrm{arctanh}(1/\sqrt{2})$ is the boost parameter. Although the boost parameter can take any value for general boosted Kerr strings, it is restricted to this value for the thin limit of a black ring. The event horizon is located at $r=r_+$ where $r_\pm:=M_K\pm\sqrt{M_K^2-a^2}$.

In studying the evaporation of a black ring, we use this boosted Kerr string solution in the following sense. We consider the situation where λ is very small, and do not take the exact limit. Then, in the neighborhood of the black ring, the spacetime metric can be well approximated by the boosted Kerr string solution. For this reason, the value of R is not infinite in our analysis although it is very large compared to M_K.

In this thin-limit approximation, the physical quantities in Eqs. (8)–(11) are expressed in terms of R, M_K and a (or a_*) as:

$$M\simeq 3\sqrt{2}\pi RM_K,\quad J_\psi\simeq 2\sqrt{2}\pi R^2M_K,\quad J_\phi\simeq 4\pi a_* RM_K^2.\tag{16}$$

$$\Omega_\phi\simeq\frac{a}{2M_K r_+}\frac{1}{\cosh\sigma},\quad \Omega_\psi\simeq\frac{1}{2R},\tag{17}$$

$$A_H\simeq 16\pi^2 r_+M_KR,\quad \kappa\simeq\frac{r_+-r_-}{4M_Kr_+}\frac{1}{\cosh\sigma}.\tag{18}$$

The inverse relations of Eq. (16) are

$$R\simeq\frac{3J_\psi}{2M},\quad M_K\simeq\frac{\sqrt{2}}{9\pi}\frac{M^2}{J_\psi},\quad a_*\simeq\frac{27\pi}{4}\frac{J_\psi J_\phi}{M^3},\tag{19}$$

and λ is expressed as

$$\lambda\simeq\frac{\sqrt{2}M_K}{R}\simeq\frac{4}{27\pi}\frac{M^3}{J_\psi^2}.\tag{20}$$

Because $\lambda^{-1/2}$ is proportional to the angular momentum J_ψ normalized by mass M, λ can be interpreted as the nondimensional rotation parameter along S^1.

Note that Ω_ϕ in Eq. (17) agrees with the angular velocity defined by the Killing generator ξ of the horizon of the boosted Kerr string[1],

$$\xi = \partial_t + \Omega_\phi \partial_\phi + V \partial_z \tag{21}$$

with

$$V = \tanh \sigma, \tag{22}$$

and κ in Eq. (18) is in agreement with the surface gravity of the horizon calculated with ξ. For a later convenience, it is useful to compare Ω_ϕ and κ with the angular velocity and the surface gravity of the horizon of an unboosted Kerr string. In the following, the quantities in the unboosted system is indicated by prime ($'$). In the unboosted system, the Killing generator of the horizon is $\xi' = \partial_{t'} + \Omega'_\phi \partial_\phi$, and Ω'_ϕ and κ' calculated from ξ' are

$$\Omega'_\phi = \frac{a}{2M_K r_+}, \qquad \kappa' = \frac{r_+ - r_-}{4M_K r_+}. \tag{23}$$

There are differences in the quantities of the boosted and unboosted systems by a factor of $1/\cosh\sigma$. This is understood as the effect of time delay in the Lorentz boost.

III. FORMULATION

In this section, we formulate the emission rates of mass and angular momenta of a black ring via Hawking radiation.

A. Emission rate

In this subsection, we derive general formulas to determine the evolution of a thin Pomeransky-Sen'kov black ring by the Hawking radiation utilizing the black string approximation. The evolution of a scalar field is governed by the Klein-Gordon equation in curved spacetime

$$(-g)^{-1/2}\, \partial_\mu \left(\sqrt{-g} g^{\mu\nu} \partial_\nu \Phi \right) = 0, \tag{24}$$

where g is the determinant of its metric.

---

[1] Our expression of Ω_ϕ does not agree with that of Ref. [19] because the definition is different.

To quantize the field, we need to expand it in terms of eigenmodes for Φ, which can be written in the black ring background as

$$\Phi = e^{-i\omega t} e^{im\phi} e^{in\psi} \Psi(x, y), \tag{25}$$

where ω, m and n are the eigenvalues for the Killing vector fields ∂_t, ∂_ψ and ∂_ϕ, respectively. By inserting this expression into Eq. (24), we obtain a second-order elliptic equation for $\Psi(x, y)$ in the (x, y) plane. This equation has a discrete series of regular solutions labeled by an integer ℓ. In the Schwarzschild string limit with $J_\phi = 0$, this series of solutions become proportional to the associate Legendre functions $P_\ell^m(x)$. Thus, the mode functions are labeled by the four parameters (ω, ℓ, m, n) in which ℓ, m and n take integer values.

In the case of a nonrotating black hole, the expected number of particles emitted per unit time is proportional to black body radiation:

$$\langle N_s \rangle \propto \frac{1}{e^{\omega_s/T_s} - 1}, \tag{26}$$

where ω_s is the energy of a scalar field in the background of the nonrotating black hole and T_s is the temperature of the horizon. Here, the temperature is defined as $T_s = \kappa_s/2\pi$, where κ_s is the surface gravity of its horizon. The temperatures appearing later are defined in a similar way.

In the rotating case, we have to replace ω_s by the energy of the mode with respect to the null geodesic generator of the black hole horizons because the mode function behaves as $\exp(-i\omega_* u_\pm)$ in the coordinates that are regular around the black hole horizon, where $u_\pm$ is the advanced time/retarded time around the horizon. In general, this null geodesic generator ξ can be written as $\xi = \partial_t + \sum_j \Omega_j \partial_{\phi^j}$ in terms of the time translation killing vector ∂_t and the rotational Killing vectors ∂_{ϕ^j}. Hence, ω_* for the mode $\propto \exp(-i\omega t + i \sum_j m_j \phi^j)$ is expressed as

$$\omega_* = \omega - \sum_j m^j \Omega_j. \tag{27}$$

Hence, the expected number of particles emitted per unit time from the black ring is given by

$$\langle N_{\rm BR} \rangle = \frac{\Gamma^{(\rm BR)}_{\ell mn}(\omega)}{e^{(\omega - n\Omega_\psi - m\Omega_\phi)/T_{\rm BR}} - 1}, \tag{28}$$

where $T_{\rm BR}$ is the temperature of the horizon and $\Gamma^{(\rm BR)}_{\ell mn}(\omega)$ is the greybody factor, which is identical to the absorption probability of the incoming wave of the corresponding modes.

This leads to the emission rates of the total mass and angular momenta, which give changes of the black ring mass M and angular momenta J_ψ and J_ϕ as

$$-\frac{d}{dt}\begin{pmatrix} M \\ J_\psi \\ J_\phi \end{pmatrix} = \frac{1}{2\pi}\sum_{\ell,m,n}\int_0^\infty d\omega \langle N\rangle \begin{pmatrix} \omega \\ n \\ m \end{pmatrix}, \tag{29}$$

where $\sum$ means summing up over all modes. Note that in this expression, it is difficult to estimate the greybody factor generally because no one has succeeded in separating the coordinates x and y, and two-dimensional numerical calculation of eigenfunctions is required.

Here, we consider the situation where the mode functions can be approximately evaluated: a black string limit discussed above. For the boosted black string (15), we can separate the wave equation, and therefore, we approximate the evolution of a scalar field in a black ring spacetime by that in a boosted Kerr string spacetime. In this situation, the scalar field can be separated as

$$\Phi = e^{-i\omega t}R(r)e^{-ikz}e^{im\phi}S_\ell^m(\theta), \tag{30}$$

where $S_\ell^m(\theta)$ is the spheroidal harmonic function. From the coordinate transformation (14), the relation between n of a black ring and k of a boosted black string is $n = 2kR\tanh\sigma$. The expected number of particles emitted per unit time is given by

$$\langle N\rangle = \frac{\Gamma_{\ell mn}(\omega)}{e^{(\omega - kV - m\Omega_\phi)/T} - 1}, \tag{31}$$

where $T = \kappa/2\pi$ is the temperature of the horizon with κ in Eq. (18), and V is the linear velocity of the horizon along the string direction, Eq. (22). $\Gamma_{\ell mn}(\omega)$ is the greybody factor of the boosted Kerr string spacetime. The emission rates (29) is evaluated by using $\langle N\rangle$ instead of $\langle N_{\rm BR}\rangle$.

B. Simplification

In the following, we simplify the expression of the emission rates (29). We normalize all quantities by the mass density M_K,

$$\tilde{\omega} = M_K\omega, \quad \tilde{k} = M_K k, \quad \tilde{\Omega}_\phi = M_K\Omega_\phi, \quad \tilde{T} = M_K T. \tag{32}$$

The normalized quantities $\tilde{\Omega}'_\phi$ and $\tilde{T}'$ in the unboosted system are introduced in the same manner. It is important to point out that the normalized angular velocity $\tilde{\Omega}_\phi$ and the normalized temperature $\tilde{T}$ depend only on a_*. We change the order of summations over ℓ and m by

$$\sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} = \sum_{m=-\infty}^{\infty} \sum_{\ell=|m|}^{\infty}, \tag{33}$$

and define

$$g^{(m)}(\tilde{\omega}, \tilde{k}) := \sum_{\ell=|m|}^{\infty} \Gamma_{\ell mn}. \tag{34}$$

Then, the emission rates can be written as

$$-\frac{d}{dt}\begin{pmatrix} M \\ J_\psi \\ J_\phi \end{pmatrix} = \frac{1}{2\pi M_K} \sum_{m=-\infty}^{\infty} \sum_{n=-\infty}^{\infty} \int_{|\tilde{k}|}^{\infty} d\tilde{\omega} \frac{g^{(m)}(\tilde{\omega}, \tilde{k})}{e^{(\tilde{\omega} - \tilde{k}V - m\tilde{\Omega}_\phi)/\tilde{T}} - 1} \begin{pmatrix} \tilde{\omega}/M_K \\ n \\ m \end{pmatrix}. \tag{35}$$

Here, the lower limit of the integral is $|\tilde{k}|$ because the mode with its energy $\omega < |\tilde{k}|$ is gravitationally bounded and does not escape to infinity. Because the number of the mode n per unit $\tilde{k}$ is very large, $O(1/\lambda)$, the summation over n can be replaced by the integral:

$$\sum_n \to \int dn = \frac{2R \tanh\sigma}{M_K} \int d\tilde{k}. \tag{36}$$

Because the integrand of Eq. (35) is zero on the boundary of the integration domain, the difference of the integral from the original discrete summation is $O(\lambda^2)$ and thus negligible. Rewriting M_K and R by M, J_ψ and J_ϕ using Eq. (19), we obtain

$$-\frac{dM}{dt} = \frac{3^7\pi^2}{2^3} \frac{J_\psi^4}{M^7} \sum_{m=-\infty}^{\infty} \int_{-\infty}^{\infty} d\tilde{k} \int_{|\tilde{k}|}^{\infty} d\tilde{\omega} \frac{\tilde{\omega} g^{(m)}(\tilde{\omega}', \tilde{k}')}{e^{(\tilde{\omega} - \tilde{k}V_z - m\tilde{\Omega}_\phi)/\tilde{T}} - 1}, \tag{37}$$

$$-\frac{dJ_\psi}{dt} = \frac{3^8\pi^2}{2^3\sqrt{2}} \frac{J_\psi^5}{M^8} \sum_{m=-\infty}^{\infty} \int_{-\infty}^{\infty} d\tilde{k} \int_{|\tilde{k}|}^{\infty} d\tilde{\omega} \frac{\tilde{k} g^{(m)}(\tilde{\omega}', \tilde{k}')}{e^{(\tilde{\omega} - \tilde{k}V_z - m\tilde{\Omega}_\phi)/\tilde{T}} - 1}. \tag{38}$$

$$-\frac{dJ_\phi}{dt} = \frac{3^5\pi}{2^2\sqrt{2}} \frac{J_\psi^3}{M^5} \sum_{m=-\infty}^{\infty} \int_{-\infty}^{\infty} d\tilde{k} \int_{|\tilde{k}|}^{\infty} d\tilde{\omega} \frac{m g^{(m)}(\tilde{\omega}', \tilde{k}')}{e^{(\tilde{\omega} - \tilde{k}V_z - m\tilde{\Omega}_\phi)/\tilde{T}} - 1}. \tag{39}$$

The integral in each formula can be further simplified if we perform transformation from $(\tilde{\omega}, \tilde{k})$ to $(\tilde{\omega}', \tilde{k}')$, where

$$\tilde{\omega} = \tilde{\omega}' \cosh\sigma + \tilde{k}' \sinh\sigma, \qquad \tilde{k} = \tilde{\omega}' \sinh\sigma + \tilde{k}' \cosh\sigma. \tag{40}$$

Substituting these formula with $\cosh\sigma = \sqrt{2}$ and $\sinh\sigma = 1$, and defining

$$I_1^{(m)}(a_*) := \int_{-\infty}^{\infty} d\tilde{k}' \int_{|\tilde{k}'|}^{\infty} d\tilde{\omega}' \frac{\tilde{\omega}' g^{(m)}(\tilde{\omega}', \tilde{k}')}{e^{(\tilde{\omega}' - m\tilde{\Omega}'_\phi)/\tilde{T}'} - 1}, \tag{41}$$

$$I_2^{(m)}(a_*) := \int_{-\infty}^{\infty} d\tilde{k}' \int_{|\tilde{k}'|}^{\infty} d\tilde{\omega}' \frac{m g^{(m)}(\tilde{\omega}', \tilde{k}')}{e^{(\tilde{\omega}' - m\tilde{\Omega}'_\phi)/\tilde{T}'} - 1}, \tag{42}$$

we obtain

$$-\frac{1}{M}\frac{dM}{dt} = 2F(a_*)\frac{J_\psi^4}{M^8}, \tag{43}$$

$$-\frac{1}{J_\psi}\frac{dJ_\psi}{dt} = 3F(a_*)\frac{J_\psi^4}{M^8}, \tag{44}$$

$$-\frac{1}{J_\phi}\frac{dJ_\phi}{dt} = G(a_*)\frac{J_\psi^3}{J_\phi M^5}, \tag{45}$$

with

$$F(a_*) := \frac{3^7\pi^2}{2^3\sqrt{2}} \sum_{m=-\infty}^{\infty} I_1^{(m)}(a_*), \tag{46}$$

$$G(a_*) := \frac{3^5\pi}{2^2\sqrt{2}} \sum_{m=-\infty}^{\infty} I_2^{(m)}(a_*). \tag{47}$$

Note that we used the fact that the integral of terms proportional to $\tilde{k}'$ vanish because such terms are odd functions of $\tilde{k}'$. From Eqs. (43)–(45) and Eq. (19), the equation of a_* is derived as

$$-\frac{1}{a_*}\frac{da_*}{dt} = 3H(a_*)\frac{J_\psi^4}{M^8}, \tag{48}$$

where

$$H(a_*) := \frac{9\pi}{4}\frac{G(a_*)}{a_*} - F(a_*). \tag{49}$$

Therefore, the time evolution of a black ring in the Hawking radiation is determined by the equations for M, J_ψ and a_*, that is, Eqs. (43), (44) and (48). The remaining work is to calculate the greybody factors and obtain $F(a_*)$ and $H(a_*)$ of Eqs. (46) and (49) numerically.

C. Greybody factor

In the following, we discuss the greybody factors for massless scalar fields in a boosted Kerr string spacetime. Substituting the ansatz (30) into the Klein-Gordon equation (24) in the background of the boosted Kerr string (15), we get the following angular and radial wave equations for $S_\ell^m(\theta)$ and $R(r)$[19],

$$0 = \frac{1}{\sin\theta}\partial_\theta\left(\sin\theta\partial_\theta S_\ell^m\right) + \left[a^2\left(\omega^2 - k^2\right)\cos^2\theta - \frac{m^2}{\sin^2\theta} + \lambda_{\ell m}\right] S_\ell^m, \tag{50}$$

$$
\begin{aligned}
0 = \Delta\partial_r\left(\Delta\partial_r R\right) - \Delta\left[k^2r^2 + a^2\omega^2 - 2\omega m a\cosh\sigma + \lambda_{\ell m}\right] R \\
+ \Bigg[\left[\omega\left(r^2+a^2\right) - ma\cosh\sigma\right]^2 + 2M_K r\left(r^2+a^2\right)\cosh^2\sigma\left(\omega - k\tan\sigma\right)^2 \\
- 2M_K r\left(r^2+a^2\right)\omega^2 - m^2a^2\sinh^2\sigma + 4kmaM_K r\sinh\sigma\Bigg] R,
\end{aligned}
\tag{51}
$$

where $\lambda_{\ell m}$ is the separation constant which is determined as an eigenvalue of (50). For small $a^2\left(\omega^2 - k^2\right)$, the eigenvalues associated with the spheroidal wave functions S_ℓ^m are $\lambda_{\ell m} = \ell\left(\ell+1\right) + \mathcal{O}\left(a^2\left(\omega^2-k^2\right)\right)$ [20].

As we mentioned in Sec. III A, the greybody factor is calculated as the absorption probability of the incoming waves of the corresponding mode. With Eq. (40), the tortoise coordinate r_* and a new wave function u defined as

$$
dr_* = \frac{r^2+a^2}{\Delta}dr, \qquad R = \frac{1}{\sqrt{r^2+a^2}}u, \tag{52}
$$

the radial wave equation (51) can be rewritten as the equation of Schrödinger type:

$$
\left[\frac{d^2}{dr_*^2} + \omega'^2 - V(r)\right] u = 0, \tag{53}
$$

where $V(r)$ is the effective potential

$$
V(r) = \frac{\Delta\left\{2M_K r(r^2-2a^2) + a^2(r^2+a^2)\right\}}{(r^2+a^2)^4} + \frac{\Delta(\omega'^2a^2 + \lambda_{\ell m} + k'^2r^2) + 4m\omega' M_K ar - m^2a^2}{(r^2+a^2)^2}. \tag{54}
$$

As the boundary condition, we impose the ingoing condition at the horizon. Then, u behaves as

$$
u(r_*) \sim \begin{cases} e^{-i\omega'_+ r_*} & \text{at} \quad r \to r_+, \\ A_{in}e^{-i\omega'_\infty r_*} + A_{out}e^{i\omega'_\infty r_*} & \text{at} \quad r \to \infty. \end{cases} \tag{55}
$$

Here $\omega'_+ := \omega' - m\Omega'_\phi$ and $\omega'_\infty := \sqrt{\omega'^2 - k'^2}$. The greybody factor is written as

$$
\Gamma = 1 - |A_{out}/A_{in}|^2, \tag{56}
$$

which has to be evaluated numerically.

In this paper, we perform numerical calculations of the greybody factors in the case of the Emparan-Reall black ring, i.e., $a = 0$. In this case, S_ℓ^m becomes the spherical harmonic function and the calculation becomes much simpler compared to the case $a \neq 0$. We developed a code to calculate the greybody factors in this situation. Here we show the example of the behavior of the greybody factors for the modes of three lowest l numbers in Fig. 2. Our result is in good agreement with the analytic approximate formula for low-frequency waves in Ref. [21].



FIG. 2: Greybody factors for the modes $k = m = 0$, $\ell = 0, 1, 2$ (from left to right).

IV. EVOLUTION BY EVAPORATION

In this section, we discuss general features of time evolution of the evaporating Pomeransky-Sen'kov black ring. Then, we focus attention to the case of the Emparan-Reall black ring, and derive semianalitic solutions of the time evolution using the numerical results of the greybody factors.

A. Evolution of Pomeransky-Sen'kov black rings

Here, we discuss the general feature that does not depend on the details of the greybody factor. From Eqs. (43) and (44), the following relation can be immediately found:

$$\frac{J_\psi^2}{M^3} = \text{const.} \tag{57}$$

From Eq. (20), this means that the parameter λ is unchanged,

$$\lambda(t) \equiv \lambda(0). \tag{58}$$

Therefore, a Pomeransky-Sen'kov black ring evaporates without changing the initial value of the nondimensional rotation parameter along S^1.

Next, let us assume $a_* > 0$ and consider how to solve $a_*(t)$ and $M(t)$. Eliminating M and J_ψ from Eqs. (43) and (48) using Eq. (57), we obtain

$$\frac{d}{dt}\left(\frac{3a_* H(a_*)}{da_*/dt}\right) = 4F(a_*). \tag{59}$$

In principle, this equation can be solved at least numerically once $H(a_*)$ and $F(a_*)$ are generated. Then, from Eqs. (43) and (48), the time evolution of $M(t)$ is formally given as

$$M(t) = M(0)\exp\left[\frac{2}{3}\int_{a_*(0)}^{a_*(t)} \frac{F(a_*)}{a_* H(a_*)} da_*\right]. \tag{60}$$

We point out that the behavior of $H(a_*)$ is crucial for the evolution of a_*. This function is analogous to $h(a_*)$ of Ref. [10] where evolution of a four-dimensional Kerr black hole was investigated: The value of a_* increases (decreases) if $H(a_*)$ is negative (positive). If $H(a_*)$ crosses zero from negative to positive at some value $a_* = a_*^{(c)}$ similarly to Fig. 3 of Ref. [10], the black ring inevitably evolves to the state with $a_*^{(c)}$. Therefore, numerical calculation of $H(a_*)$ is very interesting and we are planning to study this issue. In the next subsection, as the first step, we study what happens in the case of an Emparan-Reall black ring.

B. Evolution of Emparan-Reall black rings

In the case of the Emparan-Reall black ring, $J_\phi(t) \equiv 0$ and $a_* \equiv 0$ and Eqs. (43) and (48) can be solved analytically:

$$M(t) = M(0)\left(1 - 4F(0)\frac{J_\psi(0)^4}{M(0)^8}t\right)^{1/2}, \tag{61}$$

$$J_\psi(t) = J_\psi(0)\left(1 - 4F(0)\frac{J_\psi(0)^4}{M(0)^8}t\right)^{3/4}. \tag{62}$$

Therefore, our remaining task is to determine $F(0)$ numerically.

As discussed in Sec. III B, basically we have to calculate Eqs. (34), (41), and (46), and in those equations, the summation over m was taken at the last step. But in the case of $a_* = 0$ considered here, it is better to take summation with respect to m in advance because the integrand does not depend on m. For this reason, we calculate summation of the greybody factors as

$$g(\tilde{\omega}', \tilde{k}') = \sum_{\ell=0}^{\infty}\sum_{m=-\ell}^{\ell} \Gamma'_{\ell mn} = \sum_{l=0}^{\infty}(2\ell+1)\Gamma_{\ell mn}. \tag{63}$$

Then, $F(0)$ is given by

$$F(0) = \frac{3^7\pi^2}{2^3\sqrt{2}} I_1, \qquad I_1 := \int_{-\infty}^{\infty} d\tilde{k}' \int_{|\tilde{k}'|}^{\infty} d\tilde{\omega}' \frac{\tilde{\omega}' g(\tilde{\omega}', \tilde{k}')}{e^{\tilde{\omega}'/\tilde{T}'} - 1}. \tag{64}$$

The integration of I_1 was proceeded with the Simpson's method. The domain of integration was made finite by discarding the region where the integrand is smaller than 10^{-16}. We took summation with respect to ℓ up to $\ell = 10$, because the greybody factors for $\ell > 10$ turned out to be negligible. This is because the potential wall for $\ell > 10$ is so high that it reflects almost of all waves. In this manner, $F(0)$ is determined as

$$F(0) \simeq 0.239. \tag{65}$$

As a check, we compute $F(0)$ using the DeWitt approximation [22] in Appendix A. The two results agree well, and therefore, our numerical result is reliable.

As shown in Eq. (58), the nondimensional parameter λ is held fixed throughout the evolution. Because $a_* \equiv 0$, the constancy of the nondimensional parameter also indicates that the Emparan-Reall black ring evaporates keeping similarity to its initial shape: The black ring at any time can be obtained by uniformly scaling its initial configuration. The scaling factor can be found by deriving the time evolution of the ring radius R as

$$\frac{R}{R(0)} = \left(1 - 4F(0)\frac{J_\psi(0)^4}{M(0)^8}t\right)^{1/4}. \tag{66}$$

The lifetime T of a thin black ring with mass M is

$$T \approx \left(\frac{27\pi\lambda}{4}\right)^2 \left(\frac{M}{M_p}\right)^2 t_p, \tag{67}$$

where M_p and t_p are the Planck mass and Planck time, respectively. The time scale is proportional to M^2, and this dependence on M is same as that of the five-dimensional Schwarzschild black hole. However, because of the prefactor $(27\pi\lambda/4)^2$, the life time of a thin black ring with $\lambda \lesssim 10^{-2}$ is much shorter than the five-dimensional Schwarzschild black hole with the same mass.

V. SUMMARY

In this paper, we have studied the time evolution of evaporation of a thin black ring under the assumption that only massless scalar fields are emitted in the Hawking radiation. In order to separate the Klein-Gordon equation in the background of a black ring metric, we have considered the thin-limit approximation, where the black ring metric is approximated by the boosted Kerr string metric. Then, we have given a set of equations, Eqs. (43), (44) and (48), that determines the quasistationary evaporation of a Pomeransky-Sen'kov black ring. In this setup, a black ring evaporates without changing the nondimensional rotation parameter λ along the S^1 direction. Also, we have analytically solved these equations in the case of an Emparan-Reall black ring, Eqs. (61) and (62), with the factor $F(0) \simeq 0.239$ that has been determined by numerical calculation of the greybody factor. In the evaporation, the shape of the Emparan-Reall black ring keeps similarity to its initial configuration.

As a future work, we plan to study the time evolution of a Pomeransky-Sen'kov black ring with nonvanishing rotational parameter a_* along the S^2 direction. For this purpose, we have to calculate the functions $F(a_*)$ and $H(a_*)$ of Eqs. (46) and (49), and therefore, developing the code for computing the greybody factors of the boosted Kerr string is required.

Acknowledgments

This work was supported by the Grant-in-Aid for Scientific Research (A) (22244030).

Appendix A: DeWitt approximation

In order to check the validity of our numerical result (65) of $F(0)$ in the case of the Emparan-Reall black ring, we compute this value in an approximate way. As the method of approximation, we adopt the DeWitt approximation [22] that was originally developed to evaluate the contribution of the greybody factor to the evaporation of a Schwarzschild black hole (see p. 394 of Ref. [23] for a review). In that study, the greybody factor was obtained by appropriately reinterpreting the capture condition of null geodesics. Although this approximation holds only for high-frequency regime in a strict sense, it gives a rather good result. In fact, the difference of the DeWitt approximation from the numerical result is $\approx 6\%$. Compare the formula of the mass-loss rate by the DeWitt approximation (Eq. (146) of Ref. [22]) and the numerical value reported in Ref. [10].

In the spacetime of a five-dimensional Schwarzschild string, a massless particle with momentum in the z direction effectively behaves as a massive particle in a four-dimensional Schwarzschild spacetime. Therefore, as the first step, we study timelike geodesics in a four-dimensional Schwarzschild spacetime and derive the capture condition. Then, we translate it to the greybody factor for a massless scalar field in a spacetime of a Schwarzschild string. Using this result, we derive the value of $F(0)$ in the DeWitt approximation by performing the integration and summation in Eqs. (34), (41), and (46).

The metric of a four-dimensional Schwarzschild spacetime is given by

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2(d\theta^2 + \sin^2\theta d\phi^2), \tag{A1}$$

$$f(r) = 1 - \frac{2M_K}{r}. \tag{A2}$$

The geodesic motion of a massive particle in the equatorial plane is governed by the following equations:

$$f(r)\dot{t} = e, \tag{A3}$$

$$r^2\dot{\phi} = l, \tag{A4}$$

$$-f(r)\dot{t}^2 + \frac{\dot{r}^2}{f(r)} + r^2\dot{\phi}^2 = -1. \tag{A5}$$

Here, e and l indicate the energy and angular momentum per unit mass of the test particle, and dot ($\dot{}$) denotes the derivative with respect to particle's proper time τ. Substituting Eqs. (A3) and (A4) into Eq. (A5), we obtain

$$\dot{r}^2 + V(r) = e^2, \tag{A6}$$

where

$$V(r) = \left(\frac{l^2}{r^2} + 1\right)\left(1 - \frac{2M_K}{r}\right). \tag{A7}$$

Let us consider the situation where a test particle exists in the neighborhood of the horizon and moves toward outside (i.e. $\dot{r} > 0$). Denoting the peak value of $V(r)$ as $V_{\rm peak}$, the particle escapes to infinity if the condition $V_{\rm peak} < e^2$ is satisfied. Conversely, a particle with $V_{\rm peak} > e^2$ is reflected back to the black hole by the centrifugal barrier. After some calculation, the condition $V_{\rm peak} = e^2$ is shown to be equivalent to

$$l = e\sqrt{F(e)}M_K, \tag{A8}$$

where

$$F(e) = \frac{27e^4 - 36e^2 + 8 + e(9e^2 - 8)^{3/2}}{2e^2(e^2 - 1)}. \tag{A9}$$

Therefore, a particle escapes to infinity if $l < e\sqrt{F(e)}M_K$, and it is reflected back to the black hole if $l > e\sqrt{F(e)}M_K$. This condition is equivalent to the one obtained in Ref. [24]. Here, $\sqrt{F(e)}$ varies from 4 to $3\sqrt{3}$ as e is increased from unity to infinity.

We use this result in order to approximate the greybody factor in the particle emission from the Schwarzschild string. Here, we choose the unboosted frame, and as done in

Sec. III B, the quantities in this frame are indicated by prime ($'$). Consider the emission of a quantum particle with mass k', angular frequency ω', and angular quantum number ℓ. Replacing as $l \to \ell/k'$ and $e \to \omega'/k'$ in the capture condition derived above, the particle with $\ell \lesssim \omega'\sqrt{F(\omega'/k')}M_K$ escapes to infinity and that with $\ell \gtrsim \omega'\sqrt{F(\omega'/k')}M_K$ falls back to the horizon. Therefore, the greybody factor is approximated as

$$\Gamma_{\ell mn} \approx \theta(\tilde{\omega}'\sqrt{F(\tilde{\omega}'/\tilde{k}')} - \ell), \tag{A10}$$

where $\theta(u)$ denotes the Heaviside step function, and we introduced $\tilde{\omega}' = M_K\omega'$ and $\tilde{k}' = M_K k'$ in the same manner as Eq. (32). Note that in the massless particle limit, $\tilde{k}'/\tilde{\omega}' \to 0$, Eq. (A10) becomes $\Gamma_{\ell mn} \approx \theta(3\sqrt{3}\tilde{\omega}' - \ell)$, and this agrees with the formula in the original DeWitt approximation [22].

Now, we evaluate the value of $F(0)$. The computation can be done with the formula given in Sec. IV B. The quantity $g(\tilde{\omega}', \tilde{k}')$ in Eq. (63) is

$$g(\tilde{\omega}', \tilde{k}') = \sum_{l=0}^{\infty}(2\ell+1)\theta(\tilde{\omega}'\sqrt{F(\omega'/k')} - \ell) \approx \tilde{\omega}'^2 F(\tilde{\omega}'/\tilde{k}'). \tag{A11}$$

Here, the summation over ℓ was changed by integration as done by DeWitt. Then, $F(0)$ can be calculated by substituting this formula into Eq. (64). It is convenient to introduce new variables (x, y) by $x = \tilde{k}'/\tilde{\omega}'$ and $y = \tilde{\omega}'$, and in these variables, analytic integration can be proceeded as

$$\begin{aligned} I_1 &= \int_0^{\infty} \frac{y^4 dy}{e^{8\pi y} - 1} \int_{-1}^{1} dx F(1/|x|) \\ &= \frac{\zeta(5)}{4096\pi^5}\left[88 + 33\sqrt{2}\arcsin\left(\frac{2\sqrt{2}}{3}\right) - 3\log 3\right]. \end{aligned} \tag{A12}$$

Substituting this result into Eq. (64), we obtain

$$F(0) \approx 0.224. \tag{A13}$$

This value is fairly close to our numerical value in Eq. (65): Similarly to the original DeWitt approximation for the Schwarzschild black hole, the approximate value is about 6% smaller compared to the numerical value. Therefore, the result of the DeWitt approximation supports the correctness of our numerical calculation.

---

[1] B. Carter, Phys. Rev. Lett. **26**, 331 (1971).

[2] S. W. Hawking, Commun. Math. Phys. **25**, 152 (1972) .

[3] R. C. Myers and M. J. Perry, Annals Phys. **172**, 304 (1986).

[4] R. Emparan and H. S. Reall, Living Rev. Rel. **11**, 6 (2008) [arXiv:0801.3471 [hep-th]].

[5] R. Emparan and H. S. Reall, Phys. Rev. Lett. **88**, 101101 (2002) [hep-th/0110260].

[6] A. A. Pomeransky and R. A. Sen'kov, hep-th/0612005.

[7] S. W. Hawking, Commun. Math. Phys. **43**, 199 (1975).

[8] D. N. Page, Phys. Rev. D **13**, 198 (1976).

[9] D. N. Page, Phys. Rev. D **14**, 3260 (1976) .

[10] C. M. Chambers, W. A. Hiscock and B. Taylor, Phys. Rev. Lett. **78**, 3249 (1997) [gr-qc/9703018].

[11] B. E. Taylor, C. M. Chambers and W. A. Hiscock, Phys. Rev. D **58**, 044012 (1998) [gr-qc/9801044].

[12] H. Nomura, S. Yoshida, M. Tanabe and K. -i. Maeda, Prog. Theor. Phys. **114**, 707 (2005). [hep-th/0502179].

[13] U. Miyamoto and K. Murata, Phys. Rev. D **77**, 024020 (2008) [arXiv:0705.3150 [hep-th]].

[14] B. Chen and W. He, Class. Quant. Grav. **25**, 135011 (2008) [arXiv:0705.2984 [gr-qc]].

[15] L. Zhao, Commun. Theor. Phys. **47**, 835 (2007) [hep-th/0602065].

[16] Q. -Q. Jiang, Phys. Rev. D **78**, 044009 (2008) [arXiv:0807.1358 [hep-th]].

[17] Y. Chen, K. Hong and E. Teo, Phys. Rev. D **84**, 084030 (2011) [arXiv:1108.1849 [hep-th]].

[18] H. Elvang and M. J. Rodriguez, JHEP **0804**, 045 (2008) [arXiv:0712.2425 [hep-th]].

[19] O. J. C. Dias, Phys. Rev. D **73**, 124035 (2006) [hep-th/0602064].

[20] S. A. Teukolsky, Astrophys. J. **185**, 635 (1973).

[21] E. Jung, S. Kim and D. K. Park, JHEP **0409**, 005 (2004) [hep-th/0406117].

[22] B. S. DeWitt, Phys. Rept. **19**, 295 (1975).

[23] V. P. Frolov and I. D. Novikov, *Black Hole Physics: Basic Concepts and New Developments*, (Kluwer Academic Publishers, Dordrecht and Boston, 1998).

[24] A. F. Zakharov, Soviet Astronomy **32**, 456 (1988).